

(1) Submission ID#1522829

Title: MenB/Gonococcal Vaccine to prevent colonization and disease Authors: Serena Giuntini*, Alejandro D. Bolanos, Scarlet W. Tefera and Gregory R. Moe OMVax, Inc, San Francisco, CA *Presenter

Author(s)

Serena Giuntini, PhD

R&D Director

OMVax Inc.

Gregory R. Moe, PhD

Chief Scientific Officer

OMVax

Scarlet Tefera, BS

Research Associate I

OMVax

Alejandro Bolanos, BS

Research Associate II

OMVax

Background

MenACYW conjugate vaccines are not only protective against disease but also cost effective because they also prevent colonization thus providing community immunity in addition individual protection. We have developed a vaccine based on meningococcal native outer membrane vesicles containing overexpressed, mutant Factor H binding proteins from subfamilies A and B with reduced FH binding and attenuated endotoxin (NOMV-FHbp). Previously, we had shown that a prototype vaccine elicited serum bactericidal activity (SBA) against diverse MenB subfamily B strains in infant macaques but also SBA against gonococcal strains. In this study, we investigated the ability of antibodies elicited in mice by NOMV-FHbp to mediate SBA and prevent colonization by MenB and *N. gonorrhoeae* (Ng) strains in in vitro and in vivo models.

Aim/Methods

CD1 mice or human CEACAM1/FH transgenic (Tg) mice were immunized with two or three doses of a vaccine containing NOMV-FHbp adsorbed to AIOH3. Antisera were tested for FHbp-specific IgG by ELISA, and for SBA against Ng and diverse MenB strains. The effect of NOMV-FHbp vaccine on MenB colonization was tested in Tg mice challenged with a heterologous MenB strain and Ng in an in vitro adhesion assay with human ME180 cervical cancer cells.

Results

Mice immunized with NOMV-FHbp or recombinant FHbp had similar IgG responses against subfamily A and B FHbps even though the amount of FHbp in NOMV-FHbp was 100-fold lower. NOMV-FHbp anti-sera from CD1 mice showed similar or greater SBA compared to licensed MenB vaccines against a panel of MenB strains

irrespective of matching (100%) or mismatching ($\geq 87\%$) FHbp sequences. Human FH Tg mice had higher SBA titers when immunized with the NOMV-FHbp vaccine compared to licensed MenB vaccines ($P \leq 0.01$). NOMV-FHbp decreased CFU in the nasopharynx in MenB challenged Tg mice compared to controls ($P = 0.02$). Anti-NOMV-FHbp also had SBA titers $\geq 1:5$ against Ng strains and inhibited adhesion to ME180 cervical cells compared to control ($P = 0.0071$).

Conclusions

In mice, the NOMV-FHbp vaccine was more effective than currently licensed MenB vaccines and decreased colonization in Tg and cervical cell models demonstrating a potential to provide both individual and community immunity against MenB and gonorrhea.