

(1) Submission ID#1539518

Controlled human infection with *Neisseria lactamica* in pregnant women does not elicit mother-to-infant transmission

Author(s)

Anastasia A. Theodosiou, MA MPhil MBBChir MRCP

MRC Clinical Research Training Fellow

University of Southampton

David Cleary, BSc PhD

Assistant Professor, Microbiology and Infection

University of Birmingham

Jay Laver, BSc PhD

Senior Research Fellow

University of Southampton

Adam Dale, PhD, BSc, MBChB, MRCP, FRCPath

Clinical Lecturer

University of Southampton

Debby Bogaert, MD, PhD

Professor

University of Edinburgh

Christine E. Jones, BMBS, FRCPCH, PgcertHBE, FHEA, PhD

Associate Professor

University of Southampton

Robert Read, BMedSci(physiology), MBChB, MRCP, MD, FRCP(London), FRCP(Glasgow)

Professor

University of Southampton

Background

Neisseria lactamica carriage occurs in approximately 40% of young children, and correlates inversely with *N. meningitidis* carriage and invasive disease. Intranasal inoculation with *N. lactamica* is a safe and well-characterised human challenge model in healthy adults, although it has not previously been applied in pregnancy or childhood.

Aim/Methods

Healthy pregnant women were inoculated with 10⁵ colony-forming units *N. lactamica* Y92-1009 at 36-38 weeks gestation. Maternal and infant oral, nasopharyngeal, breast milk and blood samples were obtained at birth, and at one, four and fifteen weeks post-partum. Colonisation was investigated using selective culture and whole genome sequencing (WGS). We hypothesised that inoculation would result in maternal colonisation and horizontal transmission to the infant.

Results

Twenty-one women were inoculated, and 71% (15) became colonised with *N. lactamica* Y92-1009. The inoculation strain was detected in only one infant at one time point (at birth). Conversely, *Moraxella catarrhalis* strain-sharing (confirmed by selective culture and WGS) was observed in 35% (6/17) of inoculated mother-infant pairs completing the study. Additionally, seven women already colonised with *N. lactamica* at screening were followed up but not inoculated, and *N. lactamica* strain sharing (non-Y92-1009) was observed in one mother-infant pair in this group. Of the thirty-one women screened, 41% (7/17) with a co-habiting child aged under five years were already colonised with *N. lactamica*, compared with 0% (0/14) without a co-habiting child ($p=0.009$), and oral sampling of co-habiting children at 15 weeks post-partum revealed *N. lactamica* colonisation (non-Y92-1009) in 42% (5/12). Antibiotic use was reported by 52% (11/21) inoculated and 71% (5/7) baseline-colonised women (indications unrelated to the study), which may have affected *N. lactamica* colonisation. Serum antibody and microbiome analyses are in progress.

Conclusions

Taken together, these results suggest that *N. lactamica* does not readily transmit from colonised mothers to infants, but that transmission may occur from co-habiting children to their mothers. Conversely, these data support *M. catarrhalis* transmission from mothers to infants, consistent with research highlighting *M. catarrhalis* as a common infant commensal. This is the first ever reported peripartum respiratory human challenge study, and this model could be adapted to study microbiome and immunological development using other respiratory commensals.